

Clinical Policy: Home Ventilators

Reference Number: AR.CP.MP.184

Date of Last Revision: 03/26

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

This policy describes medical necessity criteria for noninvasive and invasive home ventilators. Noninvasive ventilation (NIV) describes the administration of positive pressure to the lungs using interfaces such as, but not limited to, nasal masks, orofacial masks, full face masks, mouthpieces, nasal pillows, or helmets.^{1,2} Invasive ventilatory support describes the administration of positive pressure to the lungs through an invasive interface, such as a tracheostomy tube or endotracheal tube.¹

Note: For criteria applicable to Medicare plans, please see MC.CP.MP.184 Home Ventilators.

Policy/Criteria

- I. It is the policy of non-Medicare health plans affiliated with Arkansas Health and Wellness, Arkansas Total Care, and QualChoice that **noninvasive home ventilators are medically necessary** for the following indications:
 - A. Initial request for the first three months of noninvasive home ventilator use for restrictive thoracic disorders, all of the following^{4,5,9,10}:
 1. Documentation of a neuromuscular disease (ex. amyotrophic lateral sclerosis) or a severe thoracic cage abnormality (ex. post-thoracoplasty for tuberculosis or severe kyphoscoliosis) and one of the following taken while member/enrollee was stable (not in acute respiratory failure):
 - a. An arterial blood gas partial pressure of carbon dioxide (PaCO₂) was measured while awake and breathing room air or on prescribed oxygen with a measurement of PaCO₂ ≥ 45 mm Hg;
 - b. Sleep oximetry demonstrates O₂ saturation ≤ 88% for at least five minutes while breathing prescribed O₂;
 - c. If neuromuscular disease is present maximal inspiratory pressure is < 60 cm H₂O, or forced vital capacity is < 50% predicted;
 2. Chronic obstructive pulmonary disease (COPD) does not contribute significantly to the pulmonary limitation;
 - B. Initial request for the first three months of noninvasive home ventilator use for chronic respiratory failure due to COPD, all of the following²²:
 1. Member/enrollee exhibits hypercapnia, demonstrated by PaCO₂ ≥ 52 mmHg by arterial blood gas obtained while awake and breathing prescribed FiO₂;
 2. Sleep apnea is not the predominant cause of the hypercapnia;
Note: Formal sleep testing is not required if the medical record demonstrates that sleep apnea is not the predominant cause of hypercapnia.
 3. Member/enrollee demonstrates at least one of the following:
 - a. Requires oxygen therapy at an FiO₂ ≥ 36% or ≥ 4L nasally;
 - b. Requires ventilatory support for more than eight hours per 24-hour period;

- c. Requires a home mechanical ventilator equipped with alarms and an internal battery due to the member/enrollee being unable to effectively breathe independently for more than a few hours, and interruption of ventilatory support without alerting the member/enrollee would pose a life-threatening risk;
- C. Initial request for the first three months of noninvasive home ventilator use for obesity hypoventilation syndrome (OHS) (also known as Pickwickian syndrome), all of the following^{5,10,14}:
1. BMI \geq 30;
 2. An initial arterial blood gas PaCO₂, done while awake and breathing at baseline and the prescribed FIO₂ is \geq 45 mm Hg;
 3. Sleep-disordered hypoventilation has been documented by polysomnography and other conditions are not considered the primary cause of hypoventilation (ex. lung parenchymal or airway disease, chest wall disorder [other than mass loading from obesity], medication use, neurologic disorder, muscle weakness, or a known congenital or idiopathic central alveolar hypoventilation syndrome);
- II. It is the policy of non-Medicare Health Plans affiliated with Centene Corporation that ***continued use of noninvasive home ventilators*** after the initial certification period is **medically necessary** when meeting the following^{5,10,22}:
- A. The device is used for at least an average of four hours per 24-hour period;
 - B. Documentation supports the following:
 1. Ongoing benefits from use of the device
- III. It is the policy of non-Medicare Health Plans affiliated with Centene Corporation that ***noninvasive home ventilators for overlap syndromes*** (presence of more than one condition, such as COPD and sleep apnea) require **secondary review** by a medical director.¹⁰
- IV. It is the policy of non-Medicare Health Plans affiliated with Centene Corporation that ***initial and ongoing use of an invasive ventilator is medically necessary*** for a long-term/chronic condition or disease affecting the ability to effectively maintain an adequate respiratory status. Examples of conditions may include neuromuscular disease, thoracic restrictive disease, or chronic respiratory failure following COPD.
- V. It is the policy of non-Medicare Health Plans affiliated with Centene Corporation that a ***second or back up noninvasive or invasive ventilator is considered medically necessary*** for the following indications¹¹:
- A. A second ventilator to serve a different purpose from the first ventilator, based on medical needs (e.g., two different types of ventilators are needed for each day, such as, a negative pressure ventilator with chest shell for one indication and a positive pressure ventilator with nasal mask the rest of the day);
 - B. A back-up ventilator for one of the following:
 1. Member/enrollee is confined to a wheelchair and requires a wheel-chair mounted ventilator during the day and another ventilator of the same type for use while in bed (unable to position the wheelchair-mounted ventilator close enough to the bed for use while sleeping). Without both pieces of equipment, member/enrollee may be prone to

- medical complications, unable to achieve appropriate medical outcomes, or may not be able to use the equipment effectively;
2. Residence in remote areas with poor emergency access.

Background

The term respiratory failure refers to the inability to adequately perform the fundamental functions of respiration, delivery of oxygen to the blood stream and removal of carbon dioxide. Respiratory failure has many causes and can be acute or chronic in nature. Typically, respiratory failure initially affects the ability to effectively move oxygen into the body, also known as oxygenation failure, or to eliminate carbon dioxide, also known as ventilatory failure.^{2,15}

Routine use of noninvasive ventilation has increased over the previous two decades, and as a result, noninvasive ventilation has become an essential tool in the management of acute and chronic respiratory failure in both the home and critical care settings.¹ Noninvasive ventilation offers increased flexibility and has become a valuable treatment option for patients with acidosis in moderate to severe respiratory distress and tachypnea with increased labored breathing due to chronic obstructive pulmonary disease (COPD) exacerbation.^{1,15}

Ventilatory support is achieved through a variety of interfaces such as oronasal mask, nasal mask, nasal prongs or full-face mask and by using a variety of ventilatory modes (e.g., volume ventilation, pressure support, cuirass ventilation, bi-level positive airway pressure [BiPAP], proportional-assist ventilation [PAV], continuous positive airway pressure [CPAP]). Oxygen is delivered via tubing through a positive pressure ventilator circuit and should be heated and humidified to improve tolerance and prevent mucosal dryness, a common side effect of prolonged noninvasive ventilation. The primary goals of home noninvasive ventilation are reduction of symptoms, improvement of quality of life, reduced readmission risk and reduction of mortality.^{1,2,3}

Invasive mechanical ventilation is primarily used to facilitate the exchange of oxygen and carbon dioxide, fully or partially, in patients with respiratory failure who no longer have the capacity to breathe spontaneously or whose ventilatory needs exceed their own ability to do so adequately. It is beneficial for protecting the airway of patients with a decreased level of consciousness, upper gastrointestinal hemorrhage, emesis, or other conditions with an increased risk of aspiration in whom noninvasive ventilation is contraindicated.^{16,17}

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2025, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
E0465	Home ventilator, any type, used with invasive interface, (e.g., tracheostomy tube)
E0466	Home ventilator, any type, used with noninvasive interface, (e.g., mask, chest shell)
E0467	Home ventilator, multi-function respiratory device, also performs any or all of the additional functions of oxygen concentration, drug nebulization, aspiration, and cough stimulation, includes all accessories, components and supplies for all functions
E0468	Home ventilator, dual-function respiratory device, also performs additional function of cough stimulation, includes all accessories, components and supplies for all functions

Reviews, Revisions, and Approvals	Revision Date	Approval Date
New Policy	7/2025	7/2025
Annual review. Revision of section of I.B. including rewording of I.B.1. and 2. with no impact on criteria, addition of new I.B.3.a.-c. regarding ventilation requirements and restructuring with previous I.B.3.a.-b. becoming I.B.4.a.-b. Removed three-month specification in Criteria II. Coding and descriptions reviewed. References reviewed and updated. Reviewed by external specialist.	03/26	03/26

References

1. Soo Hoo GW. Noninvasive ventilation in adults with acute respiratory distress: a primer for the clinician. *Hosp Pract (1995)*. 2010;38(1):16-25. doi:10.3810/hp.2010.02.275
2. Hyzy RC, Choi P. Noninvasive ventilation in adults with acute respiratory failure: Practical aspects of initiation. UpToDate. www.uptodate.com. Updated June 09, 2025. Accessed January 06, 2026.
3. Heulitt, Mark J. MD, FCCM; Ranallo, Courtney MD. Breathing in america: Disease, progress, and hope. *Pediatric Critical Care Medicine* 12(3):p e159, May 2011. doi: 10.1097/PCC.0b013e3182137d7c
4. Local coverage determination: respiratory assist devices (L33800). Centers for Medicare and Medicaid Services Web site. <https://www.cms.gov/medicare-coverage-database/view/lcd.aspx?lcdid=33800&ver=26&bc=0>. Published October 01, 2015 (revised June 09, 2025). Accessed January 12, 2026.
5. National coverage determination: durable medical equipment reference list (280.1). Centers for Medicare and Medicaid Services Web site. <http://www.cms.hhs.gov/mcd/search.asp>. Published June 09, 2025. Accessed January 15, 2026.
6. Simonds AK. Home Mechanical Ventilation: An Overview. *Ann Am Thorac Soc*. 2016;13(11):2035 to 2044. doi:10.1513/AnnalsATS.201606-454FR
7. Gay PC. Nocturnal ventilator support in COPD. UpToDate. www.uptodate.com. Updated April 21, 2025. Accessed January 06, 2026.

8. Hyzy RC, McSparron JI. Noninvasive ventilation in adults with acute respiratory failure: Benefits and contraindications. UpToDate. www.uptodate.com. Updated November 21, 2025. Accessed January 06, 2026.
9. Wilson M, Wang Z, Dobler CC, et al. *Noninvasive Positive Pressure Ventilation in the Home (with addendum)*. Rockville (MD): Agency for Healthcare Research and Quality (US); April 2, 2020.
10. Department of Health and Human Services - Office of Inspector General. Sleep Management, LLC: Audit of Claims for Monthly Rental of Noninvasive Home Ventilators. <https://oig.hhs.gov/oas/reports/region4/41804066.pdf>. Published May 2021. Accessed January 14, 2026.
11. CGS Administrators, LLC. Correct Billing and Coding of Ventilators – Revised. Joint DME MAC Publication. <https://www.cgsmedicare.com/jb/pubs/news/2024/05/cope155502a.html>. Published May 09, 2024 (revised December 04, 2025). Accessed January 15, 2026.
12. Martin TJ. Noninvasive positive airway pressure therapy for the obesity hypoventilation syndrome. UpToDate. www.uptodate.com. Updated May 01, 2025. Accessed January 06, 2026.
13. Piper A, Yee B. Clinical manifestations and diagnosis of obesity hypoventilation syndrome. UpToDate. www.uptodate.com. Published July 11, 2024. Accessed January 06, 2026.
14. Mokhlesi B, Masa JF, Brozek JL, et al. Evaluation and Management of Obesity Hypoventilation Syndrome. An Official American Thoracic Society Clinical Practice Guideline. *Am J Respir Crit Care Med*. 2019;200(3):e6-e24. doi:10.1164/rccm.201905-1071ST
15. Feller-Kopman DJ, Schwartzstein RM. The evaluation and management of the nonventilated adult with acute hypercapnic respiratory failure. UpToDate. www.uptodate.com. Updated November 14, 2025. Accessed January 06, 2026.
16. Harmon EM. Acute respiratory distress syndrome (ARDS) treatment & management. Medscape. <https://emedicine.medscape.com/article/165139-treatment#d9>. Updated July 10, 2024. Accessed January 15, 2026.
17. Hyzy RC, McSparron JI. Overview of initiating invasive mechanical ventilation in adults in the intensive care unit. UpToDate. www.uptodate.com. Updated June 24, 2025. Accessed January 06, 2026.
18. Nagler J, Cheifetz IM. Noninvasive ventilation for acute and impending respiratory failure in children. UpToDate. www.uptodate.com. Updated November 07, 2024. Accessed January 06, 2026.
19. Hill NS, Kramer NR. Noninvasive ventilation in adults with chronic respiratory failure from neuromuscular and chest wall diseases: practical aspects of initiation. UpToDate. www.uptodate.com. Updated November 14, 2025. Accessed January 07, 2026.
20. Bach JR. Noninvasive ventilatory support and mechanical insufflation-exsufflation for patients with respiratory muscle dysfunction. UpToDate. www.uptodate.com. Updated October 20, 2025. Accessed January 07, 2026.
21. Hill NS, Kramer NR. Noninvasive ventilation in adults with chronic respiratory failure from neuromuscular and chest wall diseases: patient selection and alternative modes of ventilatory support. UpToDate. www.uptodate.com. Updated November 06, 2025. Accessed January 07, 2026.

22. National coverage determination: noninvasive positive pressure ventilation (NIPPV) in the home for the treatment of chronic respiratory failure (CRF) consequent to chronic obstructive pulmonary disease (COPD) (240.9). Centers for Medicare and Medicaid Services Web site. <http://www.cms.hhs.gov/mcd/search.asp>. Published June 09, 2025. Accessed January 14, 2026.

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members/enrollees. This clinical policy is not intended to recommend treatment for members/enrollees. Members/enrollees should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members/enrollees and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members/enrollees and their representatives agree to be bound by such terms and conditions by providing services to members/enrollees and/or submitting claims for payment for such services.

Note: For Medicaid members/enrollees, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

Note: For Medicare members/enrollees, to ensure consistency with the Medicare National Coverage Determinations (NCD) and Local Coverage Determinations (LCD), all applicable NCDs, LCDs, and Medicare Coverage Articles should be reviewed prior to applying the criteria set forth in this clinical policy. Refer to the CMS website at <http://www.cms.gov> for additional information.

©2019 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.