

Clinical Policy: Ropeginterferon Alfa-2b-njft (BESREMi)

Reference Number: CP.PHAR.570

Effective Date: 03.01.22

Last Review Date: 02.26

Line of Business: Commercial, HIM, Medicaid

[Coding Implications](#)[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Ropeginterferon alfa-2b-njft (BESREMi[®]) is an interferon alfa-2b.

FDA Approved Indication(s)

BESREMi[®] is indicated for the treatment of adults with polycythemia vera (PV).

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that BESREMi[®] is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria**A. Polycythemia Vera** (must meet all):

1. Diagnosis of PV;
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age $\geq$ 18 years;
4. One of the following (a, b, or c):
 - a. One of the following (i or ii):
 - i. Failure of hydroxyurea*, unless clinically significant adverse effects are experienced or all are contraindicated;^
*Prior authorization may be required for hydroxyurea
^ For Illinois HIM requests, the step therapy requirements above do not apply as of 1/1/2026 per IL HB 5395
 - ii. Request is for treatment associated with cancer for a State with regulations against step therapy in certain oncology settings (*see Appendix D*);
 - b. Member has symptomatic low risk-PV despite aspirin and phlebotomy therapy AND meets one of the following indications for cytoreductive therapy (i, ii, iii, iv, or v):
 - i. New thrombosis or disease-related major bleeding;
 - ii. Frequent phlebotomy or intolerant of phlebotomy;
 - iii. Splenomegaly;
 - iv. Progressive thrombocytosis and/or leukocytosis;
 - v. Disease-related symptoms (e.g., pruritis, night sweats, fatigue);
 - c. For use as substitute for peginterferon alfa-2a due to product unavailability (e.g., drug shortages);
5. Documentation of JAK2 V617F mutation;

6. Member meets one of the following (a or b):
 - a. For males: Documentation of hemoglobin level > 16.5 g/dL or hematocrit level of > 49% or increased red cell mass;
 - b. For females: Documentation hemoglobin level > 16 g/dL or a hematocrit level of > 48% or increased red cell mass;
7. Request meets one of the following (a or b):*
 - a. Dose does not exceed 500 mcg every 2 weeks;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration:

Medicaid/HIM – 12 months

Commercial – 6 months or duration of request, whichever is less

B. NCCN Recommended Uses (off-label) (must meet all):

1. Diagnosis of one of the following (a, b, c, or d):
 - a. Systemic mastocytosis;
 - b. Myelofibrosis;
 - c. Essential thrombocythemia;
 - d. Chronic myeloid leukemia (CML);
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age ≥ 18 years;
4. One of the following (a or b):
 - a. For use as substitute for peginterferon alfa-2a due to product unavailability (e.g., drug shortages);
 - b. For CML as initial treatment during pregnancy;
5. Dose is within FDA maximum limit for any FDA-approved indication or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

** Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration:

Medicaid/HIM – 12 months

Commercial – 6 months or duration of request, whichever is less

C. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or

2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. All Indications in Section I (must meet all):

1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving BESREMi for a covered indication and has received this medication for at least 30 days;
2. Member is responding positively to therapy;
3. If request is for a dose increase, request meets one of the following (a or b):*
 - a. For PV, one of the following (i or ii):
 - i. For members with achievement of hematological stability for at least one year while on a stable dose of BESREMi, dose does not exceed 500 mcg every 4 weeks unless medical justification supports otherwise;
 - ii. For members who have not yet achieved hematological stability, dose does not exceed 500 mcg every 2 weeks;
 - b. New dose is within FDA maximum limit for any FDA-approved indication or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration:

Medicaid/HIM – 12 months

Commercial – 6 months or duration of request, whichever is less

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

CML: chronic myeloid leukemia
 FDA: Food and Drug Administration
 NCCN: National Comprehensive Cancer Network
 PV: polycythemia vera

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
hydroxyurea (Droxia [®] , Hydrea [®])	15 to 20 mg/kg/day	20 mg/kg/day

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s):
 - Existence of, or history of severe psychiatric disorders, particularly severe depression, suicidal ideation, or suicide attempt
 - Hypersensitivity to interferon, including interferon alfa-2b, or to any component of BESREMi
 - Moderate (Child-Pugh B) or severe (Child-Pugh C) hepatic impairment
 - History or presence of active serious or untreated autoimmune disease
 - Immunosuppressed transplant recipients
- Boxed warning(s):
 - Risk of Serious Disorders: Interferon alfa products may cause or aggravate fatal or life-threatening neuropsychiatric, autoimmune, ischemic, and infectious disorders. Monitor closely and withdraw therapy with persistently severe or worsening signs or symptoms of the above disorders.

Appendix D: States with Regulations against Redirections in Cancer

State	Step Therapy Prohibited?	Notes
FL	Yes	For stage 4 metastatic cancer and associated conditions
GA	Yes	For stage 4 metastatic cancer. Redirection does not refer to review of medical necessity or clinical appropriateness
IA	Yes	For standard of care stage 4 cancer drug use, supported by peer-reviewed, evidence-based literature, and approved by FDA

State	Step Therapy Prohibited?	Notes
IN	Yes	For advanced, metastatic cancer and associated conditions
LA	Yes [‡]	For stage 4 advanced, metastatic cancer or associated conditions. [‡] Exception if clinically equivalent therapy, contains identical active ingredient(s), and proven to have same efficacy
MS	Yes	For advanced metastatic cancer and associated conditions
NV	Yes	Stage 3 and stage 4 cancer patients for a prescription drug to treat the cancer or any symptom thereof of the covered person
OH	Yes	For stage 4 metastatic cancer and associated conditions
OK	Yes	For advanced metastatic cancer and associated conditions
PA	Yes	For stage 4 advanced, metastatic cancer
TN	Yes [^]	For cancer and associated conditions [^] Exception if step therapy is for AB-rated generic equivalent, interchangeable biological product, or biosimilar product to the equivalent brand drug
TX	Yes	For stage 4 advanced, metastatic cancer and associated conditions

Appendix E: General Information

- Per NCCN,
 - Low-risk PV: age < 60 years and no prior history of thrombosis
 - High-risk PV: age ≥ 60 years and/or prior history of thrombosis
- Per NCCN, for high-risk PV patients, preferred regimens for cytoreductive therapy include hydroxyurea or ropeginterferon alfa-2b-njft.
- Per NCCN, for low-risk PV patients, ropeginterferon alfa-2b-njft is the only preferred regimen for cytoreductive therapy. Other recommended regimens include hydroxyurea or peginterferon alfa-2a.
- Per Prescribing Information, hematological parameters are stabilized when hematocrit < 45%, platelets < 400 x 10⁹/L, and leukocytes less than 10 x 10⁹/L.
- Symptoms of disease progression include fatigue, early satiety, abdominal discomfort, inactivity, problems with concentration, night sweats, pruritus, bone pain (diffuse not joint pain or arthritis), fever (> 100 F), unintentional weight loss last 6 months.
- Poor tolerance to phlebotomy is defined as recurrent episodes of post-phlebotomy syncope despite appropriate preventive interventions.

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Polycythemia vera	Starting dose: 100 mcg SC injection every 2 weeks (50 mcg if receiving hydroxyurea). Increase the dose by 50 mcg every 2 weeks until hematological parameters are stabilized (hematocrit < 45%, platelets < 400 x 10⁹/L, and leukocytes less than 10 x 10⁹/L).	500 mcg every 2 weeks

Indication	Dosing Regimen	Maximum Dose
	Maintain the two week dosing interval at which hematological stability is achieved for at least 1 year. After achievement of hematological stability for at least 1 year on a stable dose, the dosing interval may be expanded to every 4 weeks.	

VI. Product Availability

- Single-dose prefilled syringe: 500 mcg/mL
- Single-dose prefilled pen injector: 500 mcg/0.5 mL

VII. References

1. BESREMi Prescribing Information. Burlington, MA. PharmaEssentia Corporation; June 2026. Available at: <https://www.besremi.com/>. Accessed July 10, 2026.
2. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at www.nccn.org. Accessed November 25, 2025.
3. National Comprehensive Cancer Network. Myeloproliferative Neoplasms Version 2.2025. Available at: https://www.nccn.org/professionals/physician_gls/pdf/mpn.pdf. Accessed November 25, 2025.
4. National Comprehensive Cancer Network. Systemic Mastocytosis Version 1.2025. Available at: https://www.nccn.org/professionals/physician_gls/pdf/mastocytosis.pdf. Accessed: November 25, 2025.
5. ClinicalTrials.gov. Safety study of pegylated interferon alpha 2b to treat polycythemia vera (PEGINVERA). Available at: <https://clinicaltrials.gov/ct2/show/NCT01193699>. Accessed November 6, 2025.
6. Barbui T, Thiele J, Gisslinger H, et al. The 2016 WHO classification and diagnostic criteria for myeloproliferative neoplasms: document summary and in-depth discussion. *Blood Cancer J*. 2018 Feb; 8(2): 15.
7. McMullin MF, Harrison CN, Ali S, et al; BSH Committee. A guideline for the diagnosis and management of polycythemia vera. A British Society for Hematology Guideline. *British Journal Hematology*. 2019 Jan; 184(2):176-191.
8. Gisslinger H, Zagrijtschuk O, Buxhofer-Ausch V, et al. Ropeginterferon alfa-2b, a novel IFN α -2b, induces high response rates with low toxicity in patients with polycythemia vera. *Blood*. 2015 Oct 8; 126(15): 1762–1769.
9. Tefferi A and Barbui T. Polycythemia vera: 2024 update on diagnosis, risk-stratification, and management. *Am J Hematol*. 2023;98:1465-1487.
10. Marchetti M, Vannucchi AM, Griesshammer, et al. Appropriate management of polycythaemia vera with cytoreductive drug therapy: European LeukemiaNet 2021 recommendations. *Lancet Haematol*. 2022 April;9(4):e301-e311. doi: 10.1016/S2352-3026(22)00046-1.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
C9399	Unclassified drugs or biologics
J9999	Not otherwise classified, antineoplastic drugs

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created.	11.30.21	02.22
Revised initial criteria from “JAK2V617K” to “JAK2V617F” to reflect correct mutation studied in population.	10.19.22	
1Q 2023 annual review: corrected the polycythemia vera hemoglobin and hematocrit criteria to read “>” the minimum values for men and women hemoglobin and hematocrit per the WHO diagnostic criteria; for continued therapy, added criteria that for members with achievement of hematological stability for at least one year while on a stable dose of BESREMi, dose does not exceed 500 mcg every 4 weeks unless medical justification supports otherwise; added definition of hematological stability in Appendix D per PI; references reviewed and updated.	11.02.22	02.23
1Q 2024 annual review: no significant changes; for Appendix D, added Besremi as preferred regimen for cytoreductive therapy for high risk PCV; added HCPCS codes [C9399, J9999]; references reviewed and updated.	10.13.23	02.24
Removed peginterferon alfa-2a as therapeutic alternative as no longer a preferred cytoreductive therapy for high-risk PV per NCCN; Added option for usage in low-risk PV with indications for cytoreductive therapy per NCCN; for Appendix D, added definition for low-risk and high-risk PV, removed peginterferon alfa-2a from preferred regimen for cytoreductive therapy for high-risk PV, added examples of symptoms of disease progression per NCCN.	02.13.24	05.24
1Q 2025 annual review: for Commercial line of business, added standard approval duration language “6 months or duration of request, whichever is less”; references reviewed and updated.	11.01.24	02.25
1Q 2026 annual review: added step therapy bypass for IL HIM per IL HB 5395; added bypass language for states with regulations against step therapy in certain oncology settings; extended initial approval duration from 6 to 12 months for this maintenance medication for a chronic condition; for PV, added option for usage as for use as substitute for peginterferon alfa-2a due to product unavailability per NCCN; added off-label criterion for systemic mastocytosis, myelofibrosis, essential thrombocythemia, and CML per NCCN; references reviewed and updated.	11.06.25	02.26
Updated Appendix D to include Indiana.	03.27.26	
RT4: added new prefilled pen injector formulation; updated Appendix D with revised language for Tennessee.	07.10.26	

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members

and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2022 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.