

Preemptive policy: This is a P&T approved policy and can be used after the drug is FDA approved until it is superseded by an updated policy



Clinical Policy: Povetacept (ALPN-303)

Reference Number: CP.PHAR.803

Effective Date: **FDA Approval Date**

Last Review Date: 11.26

Line of Business: Commercial, HIM/ICHRA, Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Povetacept (APLN-303^{®/TM}) is a dual inhibitor of the B-cell activating factor (BAFF) and a proliferation inducing ligand (APRIL) cytokines.

FDA Approved Indication(s) **[Pending]**

APLN-303 is indicated to reduce proteinuria in adults with primary immunoglobulin A nephropathy (IgAN) who are at risk for disease progression.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that ALPN-303 is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria*

**Criteria will mirror the clinical information from the prescribing information once FDA-approved*

A. Immunoglobulin A Nephropathy (must meet all):

1. Diagnosis of IgAN confirmed via kidney biopsy;*
2. Prescribed by or in consultation with a nephrologist;
3. Age ≥ 18 years;*
4. Documentation of both of the following (a and b):
 - a. Proteinuria of ≥ 0.5 g/day or urine protein-to-creatinine ratio (UPCR) ≥ 0.75 g/g;
 - b. Estimated glomerular filtration rate (eGFR) ≥ 30 mL/min/1.73 m²;
5. Member meets both of the following, unless contraindicated or clinically significant adverse effects are experienced (a and b, *see Appendix D*):*

**For Illinois HIM requests, the step therapy requirements below do not apply per IL HB 5395*

- a. Failure of a renin-angiotensin-aldosterone system (RAAS) inhibitor (e.g., irbesartan, losartan, lisinopril, benazepril) for at least 12 weeks;
- b. RAAS inhibitor therapy dose was at least 50% of maximum labeled dose;
6. Failure of a 12-week trial of a sodium-glucose cotransporter-2 (SGLT2) inhibitor (e.g., empagliflozin, dapagliflozin) at up to maximally indicated doses, unless clinically significant adverse effects are experienced or all are contraindicated;*
- *For Illinois HIM requests, the step therapy requirements above do not apply per IL HB 5395*
7. Failure of Filspari[®] or Vanrafia[™] at up to maximally indicated doses, unless clinically significant adverse effects are experienced or both are contraindicated;*

**For Illinois HIM requests, the step therapy requirements above do not apply per IL HB 5395*

8. ALPN-303 is not prescribed concurrently with Fabhalta[®], Voyxact[®], or Trutakna[™];
9. Dose does not exceed 80 mg every 4 weeks.*

Approval duration: 6 months

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace/ICHRA) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace/ICHRA, and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace/ICHRA) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace/ICHRA, and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace/ICHRA, and CP.PMN.53 for Medicaid.

II. Continued Therapy*

**Criteria will mirror the clinical information from the prescribing information once FDA-approved*

A. Immunoglobulin A Nephropathy (must meet all):

1. Member meets one of the following (a or b):
 - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
 - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (*refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B*);
2. Member is responding positively to therapy as evidenced by one of the following (a or b):
 - a. Decrease in UPCR from baseline;
 - b. Reduction of proteinuria as evidence by a lower total urine protein per day from baseline;
3. ALPN-303 is not prescribed concurrently with Fabhalta, Voyxact, or Trutakna;
4. If request is for a dose increase, new dose does not exceed 80 mg every 4 weeks.*

Approval duration:

Medicaid/HIM/ICHRA – 12 months

Commercial – 6 months or to the member's renewal date, whichever is longer

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):

- a. For drugs on the formulary (commercial, health insurance marketplace/ICHRA) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace/ICHRA, and CP.PMN.255 for Medicaid; or
- b. For drugs NOT on the formulary (commercial, health insurance marketplace/ICHRA) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace/ICHRA, and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace/ICHRA, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace/ICHRA, and CP.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

ACEI: angiotensin-converting-enzyme inhibitor

APRIL: a proliferation inducing ligand

ARB: angiotensin receptor blocker

BAFF: B-cell activating factor

eGFR: estimated glomerular filtration rate

FDA: Food and Drug Administration

IgAN: immunoglobulin A nephropathy

RAAS: renin-angiotensin-aldosterone system

SGLT2: sodium-glucose cotransporter-2

UPCR: urine protein-to-creatinine ratio

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Filspari (sparsentan)	400 mg PO QD	400 mg/day
Vanrafia (atrasentan)	0.75 mg PO QD	0.75 mg/day
RAAS inhibitors		
Angiotensin-converting enzyme (ACE) inhibitors (e.g., benazepril, captopril, enalapril, lisinopril, quinapril)	Varies	Varies
Angiotensin-receptor blockers (ARB) (e.g., candesartan, irbesartan, losartan, olmesartan, telmisartan, valsartan)	Varies	Varies
SGLT2 Inhibitors		

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
dapagliflozin (Farxiga [®])	10 mg PO daily	10 mg/day
Jardiance [®] (empagliflozin)	10 mg PO daily	10 mg/day

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings [Pending]

- Contraindication(s): pending
- Boxed warning(s): pending

Appendix D: General Information

- The 2025 Kidney Disease Improving Global Outcomes (KDIGO) guideline recommends initial therapy with a RAAS inhibitor (ACEI or ARB) for all patients, except patients with contraindications such as low blood pressure, bilateral renal artery stenosis, or hyperkalemia, especially due to advanced chronic kidney disease (CKD).
- The 2025 KDIGO guideline recommends initial therapy with an SGLT2 inhibitor, for patients with proteinuria > 0.5 g per day, regardless of whether the patient has hypertension.
- Patients with IgAN who are considered high risk for progressive CKD despite maximum supportive care (defined as blood pressure control, reduction of proteinuria, and lifestyle modifications) may consider treatment with corticosteroids or immunosuppressive drugs; however, there is current uncertainty over the safety and efficacy of existing immunosuppressive treatment choices. For all patients in whom immunosuppression is being considered, a detailed discussion of the risks and benefits of each drug should be undertaken with the patient recognizing that adverse treatment effects are more likely in patients with eGFR < 50 mL/min/1.73 m².

V. Dosage and Administration [Pending]

Indication	Dosing Regimen	Maximum Dose
IgAN*	80 mg SC every 4 weeks*	80 mg/month*

VI. Product Availability [Pending]

Prefilled syringe: 80 mg*

VII. References

- ALPN-303 Prescribing Information.
- ClinicalTrials.gov. Evaluation of efficacy of povetacicept in adults with immunoglobulin A nephropathy (IgAN). Available at: <https://clinicaltrials.gov/study/NCT06564142>. Accessed June 30, 2026.
- Rovin BH, Barratt J, Cook HT, et al; KDIGO IgAN and IgAV Work Group. KDIGO 2025 clinical practice guideline for the management of immunoglobulin A nephropathy (IgAN) and immunoglobulin A vasculitis (IgAV). *Kidney Int.* 2025;108(4 Suppl):S1-S71. doi:10.1016/j.kint.2025.04.004

Coding Implications [Pending]

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
Pending	Pending

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created pre-emptively	08.25.26	11.26

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to

recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2026 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.