Clinical Policy: Lutetium Lu 177 Dotatate (Lutathera, Bexlutry)

Reference Number: CP.PHAR.384
Effective Date: 05.22.18
Last Review Date: 08.26
Line of Business: Commercial, HIM/ICHRA, Medicaid

Coding Implications
Revision Log

See Important Reminder at the end of this policy for important regulatory and legal information.

Description

Lutetium Lu 177 dotatate (Lutathera®, Bexlutry™) is a radiolabeled somatostatin analog.

FDA Approved Indication(s)

Lutathera and Bexlutry are indicated for the treatment of adult (*Lutathera and Bexlutry*) and pediatric patients 12 years and older (*Lutathera only*) with somatostatin receptor-positive gastroenteropancreatic neuroendocrine tumors (GEP-NETs), including foregut, midgut, and hindgut NETs.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation® that Lutathera and Bexlutry are **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Neuroendocrine Tumors (must meet all)**:**

1. Diagnosis of one of the following somatostatin receptor-positive NETs (a, b, or c):
   a. Gastrointestinal tract or pancreas;
   b. Lung or thymus (off-label);
   c. Well-differentiated, grade 3 NET (off-label);
2. Prescribed by or in consultation with an oncologist;
3. Age ≥ 12 years;
4. One of the following (a, b, or c):
   a. Disease is recurrent, metastatic, locally advanced, or unresectable;
   b. For well-differentiated, grade 3 NETs only: Disease has all of the following characteristics (i, ii, and iii):
      i. Metastatic or locally advanced;
      ii. Unresectable;
      iii. Favorable biology (e.g., relatively low Ki-67 [< 55%]);
   c. Member has poorly controlled carcinoid syndrome associated with lung or thymus NET;
5. One of the following (a, b, or c):
   a. Member experienced disease progression while on a somatostatin analog (e.g., octreotide, lanreotide);*

   **For Illinois HIM requests, the step therapy requirements above do not apply as of 1/1/2026 per IL HB 5395*

b. Member has a well-differentiated, grade 3 NET;
c. Member has a gastrointestinal or pancreas NET with Ki-67 ≥ 10% and clinically significant tumor burden;

6. Dose does not exceed 7.4 GBq (200 mCi) every 8 weeks (± 1 week), up to a total of 4 doses.

Approval duration: 36 weeks (no more than 4 total doses)

B. Pheochromocytoma/Paraganglioma (off-label) (must meet all)**:**

1. Diagnosis of a somatostatin receptor-positive pheochromocytoma/paraganglioma;
2. Prescribed by or in consultation with an oncologist;
3. Disease is metastatic or locally unresectable;
4. Dose does not exceed 7.4 GBq (200 mCi) every 8 weeks (± 1 week), up to a total of 4 doses.

Approval duration: 36 weeks (no more than 4 total doses)

C. Other diagnoses/indications (must meet 1 or 2)**:**

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
   a. For drugs on the formulary (commercial, health insurance marketplace/ICHRA) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace/ICHRA, and CP.PMN.255 for Medicaid; or
   b. For drugs NOT on the formulary (commercial, health insurance marketplace/ICHRA) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace/ICHRA, and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace/ICHRA, and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. All Indications in Section I (must meet all)**:**

1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving Lutathera/Bexlutry for a covered indication;
2. Member is responding positively to therapy;
3. Member has not received ≥ 4 doses of Lutathera/Bexlutry;
4. If request is for a dose increase, new dose does not exceed 7.4 GBq (200 mCi) every 8 weeks (± 1 week), up to a total of 4 doses.

Approval duration: 36 weeks (no more than 4 total doses)

B. Other diagnoses/indications (must meet 1 or 2)**:**

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
   a. For drugs on the formulary (commercial, health insurance marketplace/ICHRA) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace/ICHRA, and CP.PMN.255 for Medicaid; or
   b. For drugs NOT on the formulary (commercial, health insurance marketplace/ICHRA) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace/ICHRA, and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace/ICHRA, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace/ICHRA, and CP.PMN.53 for Medicaid.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

CT: computed tomography
FDA: Food and Drug Administration
GEP-NET: gastroenteropancreatic neuroendocrine tumor
mCi: millicurie
NCCN: National Comprehensive Cancer Network

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
lanreotide (Somatuline® Depot)	90 – 120 mg SC every 4 weeks	120 mg/month
octreotide LAR (Sandostatin® LAR Depot)*	20 – 30 mg IM once monthly (20 mg may be used for pancreatic NETs)	30 mg/month
octreotide (Sandostatin®)	150 – 250 mcg SC TID	450 mcg/day

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.
**Off-label for the treatment of NETs (octreotide is only FDA-approved for the treatment of symptoms associated with carcinoid tumors) – NET dosing recommendations are per the NCCN guidelines*

Appendix C: Contraindications/Boxed Warnings
None reported

Appendix D: General Information
- Somatostatin receptor expression can be detected by somatostatin receptor-based imaging, which includes ^{68}Ga-dotatate PET/CT (preferred per the NCCN) and somatostatin receptor scintigraphy.
- Use of Lutathera/Bexlutry with somatostatin analogs:
  - Before initiating Lutathera/Bexlutry: Long-acting somatostatin analogs (e.g., long-acting octreotide) should be discontinued for at least 4 weeks prior to initiation of Lutathera/Bexlutry. Short-acting octreotide can be administered as needed up to 24 hours prior to initiating Lutathera/Bexlutry.
  - During Lutathera/Bexlutry: Administer long-acting octreotide 30 mg intramuscularly 4 to 24 hours after each Lutathera/Bexlutry dose and short-acting octreotide for symptomatic management.
  - Following Lutathera/Bexlutry: Continue long-acting octreotide 30 mg intramuscularly every 4 weeks after completing Lutathera/Bexlutry until disease progression or for up to 18 months following treatment initiation.

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
GEP-NET	7.4 GBq (200 mCi) IV every 8 weeks (± 1 week) for a total of 4 doses	7.4 BGq (200 mCi)/dose (4 doses)
NET of lung or thymus origin, pheochromocytoma, paraganglioma*		

**Off-label – dosing recommendations are per the NCCN guidelines*

VI. Product Availability
Single-dose vial for injection: 370 MBq/mL (10 mCi/mL)

VII. References
1. Lutathera Prescribing Information. Millburn, NJ: Advanced Accelerator Applications USA, Inc.; October 2024. Available at: https://www.lutathera.com. Accessed April 17, 2026.
2. Bexlutry Prescribing Information. Noblesville, IN: Cirum US LLC; September 2026. Available at: www.bexlutry.com. Accessed November 3, 2026.
3. National Comprehensive Cancer Network. Neuroendocrine and Adrenal Tumors. Version 1.2026. Available at: https://www.nccn.org/professionals/physician_gls/pdf/neuroendocrine.pdf. Accessed April 28, 2026.
4. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: http://www.nccn.org/professionals/drug_compendium. Accessed April 28, 2026.
5. Strosberg J, El-Haddad G, Wolin E, et al. Phase 3 trial of ^{177}Lu-dotatate for midgut neuroendocrine tumors. N Engl J Med. 2017; 376(2): 125-135.
6. Brabander T, van der Zwan WA, Teunissen JJM, et al. Long-term efficacy, survival, and safety of [^{177}Lu-DOTA^{0},Tyr^{3}]octreotate in patients with gastroenteropancreatic and bronchial neuroendocrine tumors. Clin Cancer Res. 2017; 1-8.

7. Clinical Pharmacology [database online]. Elsevier, Inc.; 2026. Available at: https://www.clinicalkey.com/pharmacology.

Coding Implications
Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
A9513	Lutetium Lu 177, dotatate, therapeutic, 1 millicurie (mCi)
A9699	Radiopharmaceutical, therapeutic, not otherwise classified
C9399	Unclassified drugs or biologicals

Reviews, Revisions, and Approvals	Date	P&T Approval Date
3Q 2022 annual review: no significant changes; references reviewed and updated.	05.03.22	08.22
Template changes applied to other diagnoses/indications.	09.22.22	
3Q 2023 annual review: per NCCN – for NET, added coverage for well-differentiated grade 3 NET and carcinoid syndrome, and for NETs other than the aforementioned two, revised required qualifiers to include recurrent or unresectable; for pheochromocytoma/paraganglioma, revised from "metastatic or locally advanced, and unresectable" to "metastatic or locally unresectable"; revised dosing in criteria, approval duration (from 32 weeks to 36 weeks), and Section V to reflect updated PI, which allows for every 8 week dosing "± 1 week"; updated Appendix D regarding concurrent SSA use per updated PI; references reviewed and updated.	04.20.23	08.23
3Q 2024 annual review: RT4: updated NET criteria to reflect newly approved pediatric expansion; references reviewed and updated.	05.08.24	08.24
3Q 2025 annual review: added option for first-line use in gastrointestinal or pancreas NET with Ki-67 ≥ 10% and clinically significant tumor burden per NCCN; references reviewed and updated. Added step therapy bypass for IL HIM per IL HB 5395.	06.24.25	08.25
3Q 2026 annual review: no significant changes; added ICHRA line of business; references reviewed and updated.	04.28.26	08.26
Added new brand Bexlutry to policy along with corresponding HCPCS codes [A9699, C9399].	11.03.26	

Important Reminder
This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted

standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:
For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2018 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.