Clinical Policy: Gedatolisib (Revtorpyk)

Reference Number: CP.PHAR.808
Effective Date: 12.01.26
Last Review Date: 11.26
Line of Business: Commercial, HIM/ICHRA, Medicaid

Coding Implications
Revision Log

See Important Reminder at the end of this policy for important regulatory and legal information.

Description

Gedatolisib (Revtorpyk™) is a kinase inhibitor.

FDA Approved Indication(s)

Revtorpyk is indicated in combination with fulvestrant, with or without palbociclib, for the treatment of adult patients with hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, locally advanced or metastatic breast cancer without a *PIK3CA*-mutation detected following progression on or after treatment with at least one line of endocrine therapy in the metastatic setting.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation® that Revtorpyk is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Breast Cancer (must meet all)**:**

1. Diagnosis of breast cancer;
2. Prescribed by or in consultation with an oncologist;
3. Age ≥ 18 years;
4. Disease has all of the following characteristics (a, b, c, and d):
   a. HR-positive (i.e., estrogen or progesterone receptor [ER/PR]-positive);
   b. HER2-negative;
   c. Locally advanced or metastatic;
   d. Absence of *PIK3CA* mutations;
5. Prescribed in combination with fulvestrant, with or without Ibrance®;
6. Disease has progressed on or after an endocrine-based therapy regimen (*see Appendix B*) in the metastatic setting;
7. If member is a premenopausal female, member has been treated with ovarian ablation/suppression (*see Appendix D*);
8. Request meets one of the following (a or b):*
   a. Both of the following (i and ii):
      i. Dose does not exceed 180 mg once weekly on Days 1, 8, and 15 of every 28-day cycle;
      ii. Dose is at least 130 mg per dose;

b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN.*

Approval duration:
Medicaid/HIM/ICHRA – 12 months
Commercial – 6 months or to the member's renewal date, whichever is longer

B. Other diagnoses/indications (must meet 1 or 2)**:**

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
   a. For drugs on the formulary (commercial, health insurance marketplace/ICHRA) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace/ICHRA, and CP.PMN.255 for Medicaid; or
   b. For drugs NOT on the formulary (commercial, health insurance marketplace/ICHRA) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace/ICHRA, and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace/ICHRA, and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. Breast Cancer (must meet all)**:**

1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving Revtorpyk for breast cancer and has received this medication for at least 30 days;
2. Member is responding positively to therapy;
3. If request is for a dose increase, request meets one of the following (a or b):*
   a. Both of the following (i and ii):
      i. New dose does not exceed 180 mg once weekly on Days 1, 8, and 15 of every 28-day cycle;
      ii. New dose is at least 130 mg per dose;
   b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN.*

Approval duration:
Medicaid/HIM/ICHRA – 12 months
Commercial – 6 months or to the member's renewal date, whichever is longer

B. Other diagnoses/indications (must meet 1 or 2)**:**

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):

a. For drugs on the formulary (commercial, health insurance marketplace/ICHRA) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace/ICHRA, and CP.PMN.255 for Medicaid; or
b. For drugs NOT on the formulary (commercial, health insurance marketplace/ICHRA) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace/ICHRA, and CP.PMN.16 for Medicaid; or

2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace/ICHRA, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace/ICHRA, and CP.PMN.53 for Medicaid, or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

ER: estrogen receptor
FDA: Food and Drug Administration
HER2: human epidermal growth factor receptor 2
HR: hormone receptor
LHRH: luteinizing hormone-releasing hormone
NCCN: National Comprehensive Cancer Network
PR: progesterone receptor

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Endocrine Therapy		
anastrozole (Arimidex®)	1 mg PO QD	1 mg/day
exemestane (Aromasin®)	25 mg PO QD	25 mg/day
Fareston® (toremifene)	60 mg PO QD	60 mg/day
fulvestrant (Faslodex®)	500 mg IM into the buttocks slowly (1 - 2 minutes per injection) as two 5 mL injections, one in each buttock, on days 1, 15, 29 and once monthly thereafter	500 mg/day
letrozole (Femara®)	2.5 mg PO QD	2.5 mg/day
tamoxifen (Nolvadex®, Soltamox®)	20 to 40 mg PO QD	40 mg/day
megestrol acetate	40 mg PO QID	160 mg/day

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Endocrine Therapy		
NCCN-recommended CDK4/6 inhibitors to be used in combination with aromatase inhibitors as first-line therapy		
Kisqali® (ribociclib)* Verzenio® (abemaciclib)* Ibrance® (palbociclib)	Varies	Varies

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.
**NCCN category 1 recommendation*

Appendix C: Contraindications/Boxed Warnings
None reported

Appendix D: General Information
- Breast cancer: Ovarian ablation may be accomplished by surgical oophorectomy or by ovarian irradiation. Ovarian suppression utilizes luteinizing hormone-releasing hormone (LHRH) agonists that result in suppression of luteinizing hormone and release of follicle stimulating hormone from pituitary and reduction in ovarian estrogen production. LHRH agonists include goserelin and leuprolide.

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Breast cancer	180 mg IV once weekly on Days 1, 8, and 15 of a 28-day cycle in combination with fulvestrant, with or without palbociclib, until disease progression or unacceptable toxicity. Permanently discontinue in patients who are unable to tolerate 130 mg/dose	See regimen

VI. Product Availability
Lyophilized powder in a single-dose vial: 180 mg

VII. References
1. Revtorpyk Prescribing Information. Minneapolis, MN: Celcuity Inc.; July 2026. Available at: https://www.revtorpyk.com. Accessed July 21, 2026.
2. National Comprehensive Cancer Network. Breast Cancer Version 5.2026. Available at: https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf. Accessed July 21, 2026.

Coding Implications
Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J3590	Unclassified biologics
C9399	Unclassified drugs or biologicals

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created. Revised max dosing criteria to 180 mg per week per PI.	10.06.26	11.26

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:
For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2026 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.